

FRAÏSSÉ LIMIT VIA FORCING

M. GOLSHANI  

Special issue dedicated to Professor Esfandiar Eslami

Article type: Research Article

(Received: 07 November 2023, Received in revised form 30 December 2023)

(Accepted: 18 January 2024, Published Online: 18 January 2024)

ABSTRACT. Given a Fraïssé class \mathcal{K} and an infinite cardinal κ , we define a forcing notion which adds a structure of size κ using elements of the class \mathcal{K} . We show that our construction extends the Fraïssé construction in the case $\kappa = \omega$.

Keywords: Fraïssé limit, Cohen forcing, Uncountable cardinals.

2020 MSC: Primary 03E05, 03E35, 03E55

1. introduction

The Fraïssé construction is a basic and important tool in model theory that allows to pass from a suitable class of finite structures to a canonical countable structures, called the Fraïssé limit of that class.

Suppose \mathcal{L} is a finite relational language and \mathcal{K} is a class of finite \mathcal{L} -structures closed under substructures and isomorphisms. It is called a Fraïssé class if it satisfies Joint Embedding Property (JEP) and Amalgamation Property (AP). A Fraïssé limit, denoted $\text{Flim}(\mathcal{K})$, of a Fraïssé class \mathcal{K} is the unique¹ countable ultrahomogeneous (every isomorphism of finitely-generated substructures extends to an automorphism of $\text{Flim}(\mathcal{K})$) structure into which every member of \mathcal{K} embeds.

Given a Fraïssé class \mathcal{K} and an infinite cardinal κ , we would like to force a structure of size κ which shares many properties with the Fraïssé construction.

We first consider, as an example, the special cases of linear orders, and then discuss the more general case. We also show that our construction gives the original Fraïssé construction in the case $\kappa = \omega$. The results of this paper are further extended in [1], [3] and [4],

2. Fraïssé limit of linear orders

Let \mathcal{K} be the class of finite linear orders and let κ be an infinite cardinal.

Definition 2.1. A condition in $\mathbb{P}_{\kappa, \mathcal{K}}$ is of the form $p = (A_p, \leq_p)$, where

- (1) A_p is a finite subset of κ .

✉ golshani.m@gmail.com, ORCID: 0000-0002-4689-1510

<https://doi.org/10.22103/jmmr.2024.22473.1535>

Publisher: Shahid Bahonar University of Kerman

How to cite: M. Golshani, *Fraïssé limit via forcing*, J. Mahani Math. Res. 2024; 13(4): 13 - 17.

¹The existence and uniqueness follows from Fraïssé's theorem, See [2].



© the Author(s)

(2) $(A_p, \leq_p) \in \mathcal{K}$, i.e., it is a finite linear order.

The order on $\mathbb{P}_{\kappa, \mathcal{K}}$ is defined in the natural way.

Definition 2.2. Suppose $p, q \in \mathbb{P}_{\kappa, \mathcal{K}}$. Then $p \leq q$ if and only if

- (1) $A_p \supseteq A_q$,
- (2) $\leq_q = \leq_p \cap (A_q \times A_q)$.

Lemma 2.3. $\mathbb{P}_{\kappa, \mathcal{K}}$ is c.c.c. (and in fact \aleph_1 -Knaster).

Proof. Suppose $\{p_\alpha = (A_\alpha, <_\alpha) : \alpha < \omega_1\} \subseteq \mathbb{P}_{\kappa, \mathcal{K}}$. By Δ -system lemma, we may assume that $(A_\alpha : \alpha < \omega_1)$ forms a Δ -system with root d . Since there are only finitely many different orders on d , so we can find $I \subseteq \omega_1$ of size \aleph_1 such that for all $\alpha < \beta$ in I , we have $<_\alpha \cap d \times d = <_\beta \cap d \times d$. But then $q_{\alpha, \beta} = (A_\alpha \cup A_\beta, <_\alpha \cup <_\beta)$ is a condition extending both of p_α and p_β . \square

It follows that forcing with $\mathbb{P}_{\kappa, \mathcal{K}}$ preserves cardinals and cofinalities. Let G be $\mathbb{P}_{\kappa, \mathcal{K}}$ -generic over V . The next lemma follows by a simple density argument.

Lemma 2.4. $\kappa = \bigcup_{p \in G} A_p$.

Let $\leq_G = \bigcup_{p \in G} \leq_p$. As G is a filter, \leq_G is well-defined and it is a linear order on κ .

Lemma 2.5. (κ, \leq_G) is κ -dense.

Proof. Suppose $\alpha <_G \beta$ and $p \in \mathbb{P}_{\kappa, \mathcal{K}}$. Suppose $p \Vdash \{\gamma \in \kappa : \alpha <_G \gamma <_G \beta\}$ has size $< \kappa$. By Lemma 2.3, we can find $I \in V$ of size less than κ such that $p \Vdash \check{I} \supseteq \{\gamma \in \kappa : \alpha <_G \gamma <_G \beta\}$. Let $q \leq p$ be such that $\alpha, \beta \in A_q$ and for some $\gamma \in A_q \setminus I$, $\alpha <_q \gamma <_q \beta$. Then $q \Vdash \gamma \notin I$ and $\alpha \lesssim_G \gamma \lesssim_G \beta$, which is a contradiction. \square

Lemma 2.6. Suppose $I \in V$ is an infinite subset of κ . Then $(I, <_G)$ is dense in $(\kappa, <_G)$.

Proof. Suppose $p \in \mathbb{P}_{\kappa, \mathcal{K}}$ and $p \Vdash \alpha \lesssim_G \beta$. Let $\gamma \in I \setminus A_p$ and let $q = (A_q, \leq_q)$, where $A_q = A_p \cup \{\gamma\}$ and $\alpha <_q \gamma <_q \beta$. Then $q \Vdash \gamma \in I$ and $\alpha \lesssim_G \gamma \lesssim_G \beta$. \square

Remark 2.7. Work in $V[G]$. Let $\overline{(\kappa, <_G)}$ denote the completion of $(\kappa, <_G)$. Then $\overline{(\kappa, <_G)} \cong (\mathbb{R}^{V[G]}, <)$, where $(\mathbb{R}^{V[G]}, <)$ denotes the set of real numbers in $V[G]$.

Lemma 2.8. Suppose $I, J \in V$ are non-empty subsets of κ and $I <_G J$ (i.e., $\alpha <_G \beta$ for all $\alpha \in I$ and $\beta \in J$). Then I and J are finite, in particular, there exists $\gamma \in \kappa$ such that $I <_G \gamma <_G J$.

As the next lemma shows, the structure $(\kappa, <_G)$ is rigid if κ is uncountable.

Lemma 2.9. ([4, Theorem 2]) Suppose κ is an uncountable cardinal. Then $(\kappa, <_G)$ is rigid.

3. Fraïssé limit—The general case

Suppose \mathcal{L} is a finite relational language and \mathcal{K} is a Fraïssé class. For any relation symbol $R \in \mathcal{L}$, let n_R denote its arity.

Definition 3.1. A condition p is in $\mathbb{P}_{\kappa, \mathcal{K}}$ if and only if

- (1) $p \in \mathcal{K}$.
- (2) A_p , the universe of the structure p , is a subset of κ .

The order on $\mathbb{P}_{\kappa, \mathcal{K}}$ is defined in the natural way.

Definition 3.2. Suppose $p, q \in \mathbb{P}_{\kappa, \mathcal{K}}$. Then $p \leq q$ if and only if

- (1) $A_p \supseteq A_q$,
- (2) $q = p \upharpoonright A_q$, i.e., for any relational symbol $R \in \mathcal{L}$, $R^q = R^p \cap A_q^{n_R}$.

Lemma 3.3. $\mathbb{P}_{\kappa, \mathcal{K}}$ is c.c.c. (and in fact \aleph_1 -Knaster).

Proof. Suppose $\{p_\alpha = (A_\alpha, <_\alpha) : \alpha < \omega_1\} \subseteq \mathbb{P}_{\kappa, \mathcal{K}}$. By Δ -system lemma, we may assume that $(A_\alpha : \alpha < \omega_1)$ forms a Δ -system with root d . Since there are only countably many \mathcal{K} -structures with universe d , so we can find $I \subseteq \omega_1$ of size \aleph_1 such that for all $\alpha < \beta$ in I , we have $p \upharpoonright d = q \upharpoonright d$. But then, using the amalgamation property, we can find $q_{\alpha, \beta} \in \mathcal{K}$ which extends both of p_α and q_β . \square

It follows that forcing with $\mathbb{P}_{\kappa, \mathcal{K}}$ preserves cardinals and cofinalities. Let G be $\mathbb{P}_{\kappa, \mathcal{K}}$ -generic over V . The next lemma follows by a simple density argument.

Lemma 3.4. $\kappa = \bigcup_{p \in G} A_p$.

For any relational symbol $R \in \mathcal{L}$ let $R^G = \bigcup_{p \in G} R^p$, where R^p is the interpretation of R in p . As G is a filter, R^G is a well-defined n_R -ary relation on κ . Consider the structure

$$\mathcal{M}_G = (\kappa, R^G)_{R \in \mathcal{L}}.$$

Then \mathcal{M}_G is an \mathcal{L} -structure with universe κ .

Lemma 3.5. Each element of \mathcal{K} embeds into \mathcal{M}_G .

Proof. Suppose $p \in \mathcal{K}$. We may assume that A_p , the universe of the structure p , is a subset of κ . Then the set

$$D = \{q \in \mathcal{K} : p \text{ embeds into } q\}$$

is dense, from which the lemma follows. \square

The next lemma can be proved by a simple density argument.

Lemma 3.6. Suppose $I \in V$ is an infinite subset of κ . Then for any $R \in \mathcal{L}$ and any $(\alpha_1, \dots, \alpha_i) \in \kappa^i, i < n_R$ the sets

$$\{(a_{i+1}, \dots, a_{n_R}) \in I^{n_R-i} : R^G(a_1, \dots, a_{n_R})\}$$

and

$$\{(a_{i+1}, \dots, a_{n_R}) \in I^{n_R-i} : \neg R^G(a_1, \dots, a_{n_R})\}$$

are infinite; and in fact they have the same size as I .

4. Connection with the original Fraïssé construction

In this section, we show that the Fraïssé construction can be obtained from the above results for the case $\kappa = \omega$.

Thus fix the language \mathcal{L} and the class \mathcal{K} as before. We may suppose that each $p \in \mathcal{K}$ has a domain subset of ω , so in particular \mathcal{K} , is countable. Let $\mathbb{P} = \mathbb{P}_{\omega, \mathcal{K}}$. We define the following dense subsets of \mathbb{P} :

- (1) $D_n = \{p \in \mathbb{P} : n \in A_p\}$, for $n < \omega$.
- (2) $D_{(\mathcal{A}, \mathcal{A}'), \mathcal{B}, f}^{\rightarrow} = \{p \in \mathbb{P} : p \upharpoonright A = \mathcal{A}, A' \subseteq A_p \text{ and } \exists q \text{ such that } q \upharpoonright B = \mathcal{B} \text{ and } f \text{ extends to some } \bar{f} : p \cong q\}$, where $\mathcal{A} = (A, \dots), \mathcal{B} = (B, \dots), \mathcal{A}' = (A', \dots) \in \mathcal{K}, \mathcal{A}$ is a substructure of \mathcal{A}' and $f : \mathcal{A} \cong \mathcal{B}$.
- (3) $D_{\mathcal{A}, (\mathcal{B}, \mathcal{B}'), f}^{\leftarrow} = \{q \in \mathbb{P} : q \upharpoonright B = \mathcal{B}, B' \subseteq A_q \text{ and } \exists p \text{ such that } p \upharpoonright A = \mathcal{A} \text{ and } f \text{ extends to some } \bar{f} : p \cong q\}$, where $\mathcal{A} = (A, \dots), \mathcal{B} = (B, \dots), \mathcal{B}' = (B', \dots) \in \mathcal{K}, \mathcal{B}$ is a substructure of \mathcal{B}' and $f : \mathcal{A} \cong \mathcal{B}$.

Note that

$$\{D_n\} \cup \{D_{(\mathcal{A}, \mathcal{A}'), \mathcal{B}, f}^{\rightarrow}\} \cup \{D_{\mathcal{A}, (\mathcal{B}, \mathcal{B}'), f}^{\leftarrow}\}$$

is countable and hence by the Rasiowa-Sikorski lemma, we can find a filter $G \subseteq \mathbb{P}$ which meets all the above dense sets. Then the resulting structure \mathcal{M}_G is isomorphic to the Fraïssé limit of the class \mathcal{K} .

5. Conclusion

We studied the Fraïssé construction using the method of forcing and showed that it gives a new approach to it. It also allowed us to extend the construction to higher cardinals, though in this case, the construction behaves differently.

6. Acknowledgement

The author's research has been supported by a grant from IPM (No. 97030417). The author would like to thank the reviewers for their thoughtful comments and efforts towards improving the manuscript.

References

- [1] Ackerman, Nathanael; Golshani, Mohammad; Mirabi, Mostafa; Cohen Generic Structures with Functions, preprint,
- [2] Hodges, Wilfrid Model theory. Encyclopedia of Mathematics and its Applications, 42. Cambridge University Press, Cambridge, 1993. xiv+772 pp. ISBN: 0-521-30442-3
- [3] Kostana, Ziemowit; Forcing-theoretic framework for the Fraïssé theory, PhD thesis.
- [4] Kostana, Ziemowit; Cohen-like first order structures, Ann. Pure Appl. Logic174(2023), no.1, Paper No. 103172, 17 pp.

MOHAMMAD GOLSHANI
ORCID NUMBER: 0000-0002-4689-1510
SCHOOL OF MATHEMATICS
INSTITUTE FOR RESEARCH IN FUNDAMENTAL SCIENCES (IPM)
P.O. BOX: 19395-5746
TEHRAN-IRAN.
Email address: golshani.m@gmail.com