

## COMMUTATORS-BASED GRAPH IN POLYGROUP

ELAHE MOHAMMAZADEH<sup>✉</sup>, S. AMIRMOFIDI<sup>✉</sup>, AND M. HAMIDI<sup>✉</sup>

*Special issue dedicated to Professor Esfandiar Eslami*

Article type: Research Article

(Received: 28 August 2023, Received in revised form 27 November 2023)

(Accepted: 18 January 2024, Published Online: 18 January 2024)

**ABSTRACT.** In this paper, first, we study commutators of a polygroup. Then for a finite polygroup  $P$  and a fixed element  $g \in P$ , we introduce the  $g$ -graph  $\Delta_P^g$ . In addition, with some additional conditions, we see that it is connected and the diameter is at most 3. Then, we investigate isomorphic graphs. Specially, we obtain a new isomorphic graph derived from an isomorphic graph and two non-commutative isomorphic polygroups. Also, we show that two polygroups with isomorphic graphs preserve nilpotency.

*Keywords:* Polygroup, Related graph, Commutator.

*2020 MSC:* Primary 20N20, 20F19, 5C25.

### 1. Introduction

Graph theory has been used to prove fundamental results in other areas of mathematics. Also, it has some applications in computer science, networking, etc. Today, modeling real-world problems based on graphs is one of the most important solutions to problems. Pure modeling without any comprehensive information about the data cannot be a proper modeling. When a set is equipped with algebraic and hyperalgebra operations or hyperoperations, it can find a regular structure or hyperstructure and act as a purposeful system. Therefore, the modeling that is based on an algebraic hyperstructure can be accurate and comprehensive and examine problem-solving in a real way. Many researches have been done in the field of graphs and algebraic structures, which have many applications. Knowledge of graphs can innovate new ideas for studying other algebraic structures such as groups and polygroups. For some extensive papers on assigning a graph to a ring, group, etc you can see [1, 8, 12, 13]. In [14], translating some properties of polygroups into graph-theoretic language, help us to study nilpotent polygroups. The theory of algebraic hyperstructures, that is a well-established branch in algebraic theory was first introduced by Marty [11]. Since then many researchers have worked on algebraic hyperstructures and developed them. The application of hyperstructures has been studied in geometry, automata, probabilities, and so on (see [5, 9]). Polygroups as a

✉ m.hamidi@pnu.ac.ir, ORCID: 0000-0002-8686-6942

<https://doi.org/10.22103/jmmr.2024.22105.1501>

Publisher: Shahid Bahonar University of Kerman

How to cite: E. Mohammadzadeh, S. Amirmofidi, M. Hamidi, *Commutators-based graph in polygroup*, J. Mahani Math. Res. 2024; 13(4): 39-52.



© the Author(s)

certain subclass of hypergroups are studied in [6]. Polygroup theory extends some well-known group results and introduces new topics such as Nilpotent and Engel polygroups (see [2, 15]). Polygroups have been discussed by Corsini (see [3, 4]), Ameri [2], Comer [5], Davvaz [6] and so on. Now, we are going to introduce a new graph called  $g$ -graph, for a fixed element  $g \in P$ . A main graph related to a polygroup is a non-commuting graph. Here, we check the conditions that the  $g$ -graph coincides with the non-commuting graph. In addition, we obtained some results on nilpotent polygroups. Also, we obtain a necessary and sufficient condition between the isomorphic graph of polygroups and groups.

## 2. Preliminaries

We recall some basic definitions which are proposed by the pioneers of this subject. A *hypergroupoid* is a non-empty set  $H$  with a *hyperoperation* “ $\circ$ ” defined on  $H$ , that is, a mapping of  $H \times H$  into the family of non-empty subsets of  $H$ . If  $(x, y) \in H \times H$ , its image under  $\circ$  is denoted by  $x \circ y$ . If  $A, B$  are non-empty subsets of  $H$ , then  $A \circ B = \bigcup \{x \circ y \mid x \in A, y \in B\}$ . For any given  $x \in H$ , we identify  $x \circ A$  for  $\{x\} \circ A$  and  $A \circ x$  for  $A \circ \{x\}$ . Generally, the singleton  $a$  is identified with its member  $a$ . The structure  $(H, \circ)$  is called a *semihypergroup* if for all  $a, b, c \in H$ ,  $a \circ (b \circ c) = (a \circ b) \circ c$ . A semihypergroup  $(H, \circ)$  is a *hypergroup*, if for any given  $x \in H$ ,  $x \circ H = H \circ x = H$ , for all  $x \in H$ .

**Definition 2.1.** [6] A polygroup is a hypergroup  $\langle P, \cdot, e, {}^{-1} \rangle$ , where  $e \in P$  and  ${}^{-1}$ , is an unitary operation on  $P$  and for all  $x, y, z \in P$ , the following axioms hold:

- (i)  $e \cdot x = x \cdot e = x$ ;
- (ii)  $x \in y \cdot z \Leftrightarrow y \in x \cdot z^{-1} \Leftrightarrow z \in y^{-1} \cdot x$ .

**Definition 2.2.** [6] A non-empty subset  $K$  of a polygroup  $P$  is a *subpolygroup* of  $P$  if for any  $a, b \in K$  we have  $ab \subseteq K$  and  $a^{-1} \in K$ . Also, a subpolygroup  $N$  of  $P$  is *normal* if for any  $a \in P$ ,  $a^{-1}Na \subseteq N$ .

A polygroup is called commutative if for any  $x, y \in P$ ,  $x \cdot y = y \cdot x$ . Let  $(P, \circ)$  and  $(P, \star)$  be two polygroups. A map  $f : (P, \circ) \rightarrow (P, \star)$  is called a *homomorphism*, if for any  $a, b \in P$ ,  $f(a \circ b) \subseteq f(a) \star f(b)$ . Also,  $f$  is a *good homomorphism*, if for any  $a, b \in P$ ,  $f(a \circ b) = f(a) \star f(b)$ . In addition,  $(P_1 \times P_2, \bigcirc)$ , where  $\bigcirc$  is defined as follows, is a polygroup (see [6]):

$$(x_1, y_1) \bigcirc (x_2, y_2) = \{(x, y) \mid x \in x_1 \circ x_2, \text{ and } y \in y_1 \star y_2\}.$$

For an equivalence relation  $\rho \subseteq P \times P$  and non-empty subsets  $A$  and  $B$  of  $P$ , it is defined  $A\bar{\rho}B \iff a\rho b$ , which,  $a \in A$  and  $b \in B$  are arbitrary. The relation  $\rho$  is called strongly regular if for all  $x, y, a \in P$ ,

$$x\bar{\rho}y \implies (a \circ x) \bar{\rho} (a \circ y) \text{ and } (x \circ a) \bar{\rho} (y \circ a).$$

**Theorem 2.3.** [6] *If  $(H, \cdot)$  is a hypergroup and  $\rho$  is a strongly regular relation on  $H$ , then  $H/\rho$  is a group under the operation  $\rho(x) \otimes \rho(y) = \rho(z)$ , for all  $z \in x \cdot y$ .*

For all  $n \geq 1$ , we define the relation  $\beta_n$  on a semihypergroup  $H$  as follows:

$$a\beta_n b \iff \exists(x_1, \dots, x_n) \in H, \text{ such that } \{a, b\} \subseteq \prod_{i=1}^n x_i$$

and  $\beta = \bigcup_{n \geq 1} \beta_n$ , where  $\beta_1 = \{(x, x) \mid x \in H\}$ , is the diagonal relation on  $H$ . This relation was introduced by Koskas [10]. Suppose that  $\beta^*$  is the transitive closure of  $\beta$ , the relation  $\beta^*$  is a strongly regular relation. Also we have:

**Theorem 2.4.** [7] *If  $H$  is hypergroup, then  $\beta = \beta^*$ .*

The kernel of the canonical map  $\varphi : P \rightarrow P/\beta^*$  is called the *core* (or heart) of  $P$  and it is denoted by  $\omega_P$  (or  $\omega$ ). It is easy to prove that  $\omega_P = \beta^*(e)$  and  $\beta^*(x)^{-1} = \beta^*(x^{-1})$  for all  $x \in P$  (see [6]).

**Theorem 2.5.** [6] *Assume  $P_1$  and  $P_2$  are two polygroups. Then  $\omega_{P_1 \times P_2} = \omega_{P_1} \times \omega_{P_2}$ .*

From now on, let  $(P, \cdot)$  is a polygroup,  $n \in \mathbb{N}$  and for any  $x \in P$ , we use  $\bar{x}$  and  $xy$  instead of  $\beta^*(x)$  and  $x \cdot y$ , respectively.

**Theorem 2.6.** [14] *For any  $x, y \in P$ ,  $\bar{x} \otimes \bar{y} = \bar{y} \otimes \bar{x}^\alpha$  if and only if  $xy\omega = yx^\alpha\omega$ .*

In Theorem 2.6, take  $\alpha$  be the identity map  $i$ . Then we get the following corollary.

**Corollary 2.7.** *Let  $x, y \in P$ . Then  $xyw = yxw$  if and only if  $\bar{x} \otimes \bar{y} = \bar{y} \otimes \bar{x}$ .*

**Definition 2.8.** [14]

(i) For an automorphism  $\alpha$  the set  $\zeta^\alpha(P)$  is called  $\alpha$ -center of  $P$  and is defined as follows:

$$\zeta^\alpha(P) = \{x \in P \mid xy\omega = yx^\alpha\omega \text{ for any } y \in P\}.$$

(ii) Center of  $P$ , denoted by  $Z(P)$ , is defined by  $\{x \in P \mid xy\omega = yx\omega \text{ for all } y \in P\}$ .

(iii) For  $x \in P$  the *centralizer*  $x$  in  $P$  is defined by  $C_P(x) = \{y \in P \mid yx\omega = xy\omega\}$ .

In [15], we see that  $Z(P)$  is a normal subpolygroup of  $P$  and  $C_P(x)$  is a subpolygroup of  $P$ .

Assume  $Z_0(P) = \omega$ , and for  $n \geq 2$ ,

$$Z_n(P) = \{x \in P \mid xyZ_{n-1}(P) = yxZ_{n-1}(P) \text{ for any } y \in P\}.$$

It is proved that  $Z_n(P)$  is a normal subpolygroup of  $P$ .

**Definition 2.9.** [15] A polygroup  $P$  is called nilpotent if and only if for some  $n \in \mathbb{N}$ ,  $Z_n(P) = P$ .

*Remark 2.10.* Clearly  $Z_n(P) = \{x \in P \mid [x, y] \subseteq \omega \text{ for any } y \in Z_{n-1}(P)\}$ .

**Theorem 2.11.** [6] Let  $(G, \cdot)$  be a group. Then  $(P_G, \circ, e, {}^{-1})$  is a polygroup, where  $P_G = G \cup \{a\}$ ,  $a \notin G$  and  $\circ$  is defined as follows:

- (i)  $a \circ a = e$ ,
- (ii)  $e \circ x = x \circ e = x$ ,  $\forall x \in G$ ,
- (iii)  $a \circ x = x \circ a = x$ ,  $\forall x \in G - \{e, a\}$ ,
- (iv)  $x \circ y = x \cdot y$ ,  $\forall (x, y) \in G^2; y \neq x^{-1}$ ,
- (v)  $x \circ x^{-1} = x^{-1} \circ x = \{e, a\}$ ,  $\forall x \in G - \{e, a\}$ .

Clearly,  $Z(P_G) = Z(G) \cup \{e, a\}$  and  $\omega_{P_G} = \{e, a\}$  and  $|P_G| = |G| + 1$ .

We recall from [17], that for a group  $(G, \cdot)$  the *commutator* of  $x, y \in G$  is  $[x, y] = x^{-1} \cdot y^{-1} \cdot x \cdot y$ , and inductively,  $[x, y_1, y_2, \dots, y_n] = [[x, y_1, y_2, \dots, y_{n-1}], y_n]$ , for any  $y_1, y_2, \dots, y_n \in G$ . Also,  $G$  is called *nilpotent* if the upper central series  $Z_n(G)$  terminates at finite steps, where

$$Z_n(G) = \{x \in G; [x, y_1, y_2, \dots, y_n] = e \text{ for any } y_1, y_2, \dots, y_n \in G\}.$$

In a graph a *path*  $\rho$  is a sequence  $v_0 e_1 v_1 \dots e_k v_k$  whose terms are alternately distinct vertices and distinct edges, such that the ends of  $e_i$  are  $v_{i-1}$  and  $v_i$  for any  $i$ ,  $1 \leq i \leq k$ . In this case,  $\rho$  is called a path between  $v_0$  and  $v_k$  and the number  $k$  is called the *length* of  $\rho$ . If  $v_0$  and  $v_k$  are adjacent by an edge  $e_{k+1}$ , then  $\rho \cup \{e_{k+1}\}$  is called a *cycle*. The *length* of a cycle define the number of its edges. The length of the shortest cycle in a graph  $\Delta$  is called *girth* of  $\Gamma$  and denoted by  $\text{girth}(\Delta)$ . If  $v$  and  $w$  are vertices of  $\Delta$ , then  $d(v, w)$  denotes the length of the shortest path between  $v$  and  $w$ . The largest distance between all pairs of the vertices of  $\Delta$  is called the *diameter* of  $\Delta$ , and is denoted by  $\text{diam}(\Delta)$ . A graph is *connected* if there is a path between each pair of the vertices of  $\Delta$ .

### 3. Commutators of a polygroup

In this section, we introduce the notion of commutator of elements in polygroups and investigate the properties of commutators of polygroup.

The commutator of two elements  $x, y \in P$  is defined by  $[x, y] = \{t \in P; t \in x^{-1}y^{-1}xy\}$ . If  $z \in P$  and  $A \subseteq P$ , then  $[A, z] = \{t; t \in A^{-1}z^{-1}Ay\}$ . Therefore

$$[[x, y], z] = \{t; t \in [x, y]^{-1}z^{-1}[x, y]z\}.$$

And inductively, we define  $[x, y_1, y_2, \dots, y_n] = [[x, y_1, y_2, \dots, y_{n-1}], y_n]$ . Also,  $A^x = \{t; t \in x^{-1}Ax\}$  (see [15]).

**Theorem 3.1.** Consider  $x, y, z, a, g \in P$  and  $A \subseteq P$ . Then

- (i)  $[xy, z] \subseteq [x, z]^y[y, z]$ . Moreover, if  $y = z$ , then  $[xy, y] = [x, y]^y$ . Also,  $[Ay, y] = [A, y]^y$ .

- (ii)  $[A, B]^{-1} = [B, A]$ .  
 (iii) If  $a \in A$ , then  $[a, y] \subseteq [A, y]$ .  
 (iv)  $[x^{-1}, y] \subseteq [x, y]^{-x^{-1}}$ .  
 (v)  $[x, yz] \subseteq [x, z][x, y]^x$ . Also,  $[x, y^{-1}] \subseteq [y, x]^{-y^{-1}}$ .

*Proof.* (i) Let  $t \in [xy, z] = \{t \in (xy)^{-1}z^{-1}xyz = y^{-1}x^{-1}z^{-1}xyz\}$ . Then  
 $t \in y^{-1}x^{-1}z^{-1}xeyz \subseteq y^{-1}x^{-1}z^{-1}xzez^{-1}yz \subseteq y^{-1}x^{-1}z^{-1}xzyy^{-1}z^{-1}yz$ .

Therefore,  $t \in mn$  for some  $m \in [x, z]^y$  and  $n \in [y, z]$ .

Thus,  $[xy, z] \subseteq [x, z]^y[y, z]$ .

Moreover,

$$\begin{aligned} t \in [x, y]^y &\Leftrightarrow t \in y^{-1}(x^{-1}y^{-1}xy)y = (xy)^{-1}y^{-1}(xy)y \\ &\Leftrightarrow t \in [xy, y]. \end{aligned}$$

Hence,  $[x, y]^y = [xy, y]$ . Also, we have

$$\begin{aligned} t \in [Ay, y] &\Leftrightarrow t \in (Ay)^{-1}y^{-1}(Ay)y \\ &\Leftrightarrow t \in y^{-1}A^{-1}y^{-1}Ayy \Leftrightarrow t \in y^{-1}[A \cdot y]y \\ &\Leftrightarrow t \in [A \cdot y]^y. \end{aligned}$$

Consequently,  $[Ay, y] = [A, y]^y$ .

Parts (ii) and (iii) are clear.

(iv) Assume  $t \in [x^{-1}, y]$ . Then

$$t \in (xy^{-1}x^{-1}ye) \subseteq xy^{-1}x^{-1}yxx^{-1},$$

and so  $t \in [x, y]^{-x^{-1}}$ .

(v) By part (i) and (ii) we have

$$\begin{aligned} [x, yz] &= [yz, x]^{-1} \subseteq ([y, x]^z[z, x])^{-1} \\ &= [z, x]^{-1}[y, x]^{-z} \\ &= [x, z][x, y]^z. \end{aligned}$$

Also, by parts (i), (ii) and (iv),  $[x, y^{-1}] = ([y^{-1}, x])^{-1} \subseteq [y, x]^{-y^{-1}}$ . Therefore,  $[x, y^{-1}] \subseteq [y, x]^{-y^{-1}}$ . This complete the proof  $\square$

In what follows, we examined that the equalities existing in group theory do not exist in polygroups, in general.

**Example 3.2.** Let  $P = \{e, a, b, c\}$  be a polygroup by the Table 1.

Then  $[ea, e] \neq [e, e]^a[a, e]$ , since

$$[ea, e] = [a, e] = ba = \{e, a, b\} \text{ and } [e, e]^a[a, e] = (ba)(ba) = \{e, a, b, e\}.$$

Also,  $[b^{-1}, a] \neq [b, a]^{-b^{-1}}$ , since  $[b^{-1}, a] = bbaa = ba = \{e, a, b\}$  and  $[b, a]^{-b^{-1}} = bbaaba = \{e, a, b, c\}$ .

**Theorem 3.3.** Let  $P$  and  $H$  be two polygroups. Then

- (i)  $Z(P \times H) = Z(P) \times Z(H)$ .

TABLE 1. Polygroup  $(P, \cdot)$ .

$\cdot$	$e$	$a$	$b$	$c$
$e$	$e$	$a$	$b$	$c$
$a$	$a$	$a$	$P$	$c$
$b$	$b$	$\{e, a, b\}$	$b$	$\{b, c\}$
$c$	$c$	$\{a, c\}$	$c$	$P$

$$(ii) (P \times H) - Z(P \times H) = (P - Z(P)) \times (H - Z(H)).$$

*Proof.* (i) Let  $(x, a) \in Z(P \times H)$ . Then  $(x, a)(y, b)\omega_{P \times H} = (y, b)(x, a)\omega_{P \times H}$  for any  $x, y \in P$  and  $a, b \in H$ . It follows by  $\omega_{P \times H} = \omega_P \times \omega_H$  that  $xy\omega_P = yx\omega_P$  and  $ab\omega_H = ba\omega_H$  and so  $x \in Z(P)$  and  $a \in Z(H)$ , Consequently,  $(x, a) \in Z(P) \times Z(H)$  i.e,  $Z(P \times H) \subseteq Z(P) \times Z(H)$ . In a similar way,  $Z(P) \times Z(H) \subseteq Z(P \times H)$ . Therefore,  $Z(P \times H) = Z(P) \times Z(H)$ .

(ii) Assume  $(x, a) \in (P \times H) - Z(P \times H)$ . By (i), we have  $(x, a) \in (P \times H) - (Z(P) \times Z(H))$  and so  $x \in P - Z(P)$  and  $a \in H - Z(H)$ . Therefore,  $(x, a) \in (P - Z(P)) \times (H - Z(H))$  i.e,  $(P \times H) - Z(P \times H) \subseteq (P - Z(P)) \times (H - Z(H))$ . Similarly,

$$(P \times H) - Z(P \times H) \supseteq (P - Z(P)) \times (H - Z(H)).$$

Consequently,  $(P \times H) - Z(P \times H) = (P - Z(P)) \times (H - Z(H))$ .  $\square$

**Theorem 3.4.** *Let  $G$  be a group. Then  $G$  is a nilpotent if and only if  $P_G$  is nilpotent.*

*Proof.* ( $\Rightarrow$ ) Assume  $G$  is a nilpotent group of class  $n$  and  $x, y_1, \dots, y_n$  are arbitrary elements of  $G$ . Then  $Z_n(G) = G$ . Thus by Theorem 2.11,  $[x, y_1, \dots, y_n] \subseteq \omega$ . On the other hand, for  $a \in P_G$ , we have  $[x, a] \subseteq \omega$ . Consequently,  $P_G$  is a nilpotent polygroup.

( $\Leftarrow$ ) Assume  $P_G$  is a nilpotent of class  $n$ . Then for any  $x, y_1, \dots, y_n \in G \subseteq P_G$ , we have  $[x, y_1, \dots, y_n] \subseteq \omega$ . By Theorem 2.11 and

$$x = \bar{x} \text{ and } y_1 = \bar{y}_1, \dots, y_n = \bar{y}_n, \text{ we have}$$

$$[x, y_1, \dots, y_n] = [x, \bar{y}_1, \dots, \bar{y}_n] = \bar{e} = \{e, a\}.$$

Moreover,  $[x, y_1, \dots, y_n] = e$  or  $a$ . But  $[x, y_1, \dots, y_n] = a$ , is a contradiction. Thus  $[x, y_1, \dots, y_n] = e$  and so  $Z_n(G) = G$ , i.e  $G$  is a nilpotent group.  $\square$

#### 4. The $g$ -graph of a polygroup

In [14], a graph  $\Gamma_P^\alpha$  is associated to a polygroup  $P$ , whose vertices are elements of  $P \setminus \zeta^\alpha(P)$  and  $x$  connected to  $y$  by edge in case  $xy\omega \neq yx^\alpha\omega$  or  $yx\omega \neq xy^\alpha\omega$ .

Now, if  $\alpha$  is the identity automorphism  $i$ , then we call  $\Gamma_P^i$  the non-commuting

graph and denoted by  $\Delta_P$ . In addition, if  $x$  connected to  $y$  by edge in case  $xy\omega = yx\omega$ , then we call it the *commuting graph and denote by  $\tau_P$* .

**Definition 4.1.** For any polygroup  $P$  and a fixed element  $g \in P$ , the  $g$ -graph of  $P$  is the graph with vertex set  $P - Z(P)$  and two distinct vertices  $x$  and  $y$  join by an edge if  $[x, y] \not\subseteq g\omega$  and  $g^{-1}\omega$  or  $[y, x] \not\subseteq g\omega$  and  $g^{-1}\omega$ .

We denote by  $V(P)$ , the set of vertices of  $\Delta_P^g$ . Also, we use  $\{x, y\}$  if there exist an edge between two vertices  $x$  and  $y$ .

**Example 4.2.** Since each polygroup of order less than 4 is commutative, so its graph is empty. Also, the graph of non-commutative polygroup  $(P = \{e, a, b, c\}, \cdot)$ , where  $\cdot$  is defined on  $P$  by the Table 2, is empty since,  $Z(P) = P$ .

TABLE 2. Polygroup  $(P, \cdot)$ .

$\cdot$	$e$	$a$	$b$	$c$
$e$	$e$	$a$	$b$	$c$
$a$	$a$	$a$	$P$	$c$
$b$	$b$	$\{e, a, b\}$	$b$	$\{b, c\}$
$c$	$c$	$\{a, c\}$	$c$	$P$

**Example 4.3.** Let  $P = \{e, a, b, c, d, f, g\}$ . We consider the proper non-commutative polygroup  $(P, \cdot, e, -1)$  ([6]), where  $\cdot$  is defined on  $P$  as Table 3.

TABLE 3. Polygroup  $(P, \cdot)$ .

$\cdot$	$e$	$a$	$b$	$c$	$d$	$f$	$g$
$e$	$e$	$a$	$b$	$c$	$d$	$f$	$g$
$a$	$a$	$e$	$b$	$c$	$d$	$f$	$g$
$b$	$b$	$b$	$\{e, a\}$	$g$	$f$	$d$	$c$
$c$	$c$	$c$	$f$	$\{e, a\}$	$g$	$b$	$d$
$d$	$d$	$d$	$g$	$f$	$\{e, a\}$	$c$	$b$
$f$	$f$	$f$	$c$	$d$	$b$	$g$	$\{e, a\}$
$g$	$g$	$g$	$d$	$b$	$c$	$\{e, a\}$	$f$

First, note that  $\omega = Z(P) = \{e, a\}$ .  $b\omega = b^{-1}\omega = \{b\}$ . In addition,  $[g, d] = g^{-1}d^{-1}gd = bc = g$ . Similarly,

$$[b, f] = [b, d] = [g, c] = [c, f] = [g, d] = g, [b, g] = [f, d][b, c] = [c, d] = f, [f, g] = \{e, a\}.$$

Therefore,  $\Delta_P^b$  is a graph with  $V(P) = \{b, c, d, f, g\}$  and  $\{b, c\}$  is an edge, because  $[b, c] = f \not\subseteq b\omega = \{b\}$ . In a same way, for any  $x, y \in V(P)$ ,  $\{x, y\}$  is an edge. Consequently,  $\Delta_P^b$  is a connected graph.

**Example 4.4.** Let  $P$  be a polygroup as Example 4.3. Then  $f\omega = \{f\}$  and  $f^{-1}\omega = \{g\}$ . Therefore,  $\Delta_P^f$  is a graph with  $V(P) = \{b, c, d, f, g\}$ . Note that  $[f, g] = \{e, a\} \not\subseteq f\omega, f^{-1}\omega$ , i.e.,  $\{f, g\}$  is an edge. But  $\{b, c\}$  is not an edge, because  $[b, c] = f \not\subseteq f^{-1}\omega = \{g\}$  and  $[b, c] = f \subseteq f\omega = \{f\}$ . Similarly,

$\{f, d\}, \{f, c\}, \{f, b\}, \{g, c\}, \{g, b\}, \{g, d\}, \{d, c\}, \{b, d\}, \{b, c\}$

are not edges. The diagram of  $\Delta_P^f$  is shown as Figure 1.




FIGURE 1. Graph  $\Delta_P^f$

*Remark 4.5.* From now on,  $K(P) = \{[x, y], x, y \in P\}$ .

*Remark 4.6.* If  $g \in \omega$  and  $\{x, y\}$  be an edge in  $\Delta_P^g$ , then  $[x, y] \not\subseteq g\omega = \omega$ , and so  $xy\omega \neq yx\omega$ . Therefore,  $\Delta_P^g$  coincides with  $\Delta_P$ . In addition, if for any  $x, y \in P$ , we have  $[x, y] \not\subseteq g\omega$  and  $g^{-1}\omega$ , then  $\Delta_P^g$  is a connected graph. Hence, in this paper, we assume  $g \notin \omega$  and there are  $x, y \in P$  such that  $[x, y] \subseteq g\omega, g^{-1}\omega$ .

**Definition 4.7** (Spanning subgraph). A graph  $\tau'$  is called spanning subgraph of  $\tau$ . We denote by  $\tau' \subseteq \tau$ , if

- (i)  $\tau'$  is a subgraph of  $\tau$ .
- (ii)  $V(\tau') = V(\tau)$ .

**Example 4.8.** Suppose  $\Delta_P^f$  and  $\Delta_P^b$  are as Examples 4.3 and 4.4. Clearly,  $\Delta_P^f \subseteq \Delta_P^b$ .

**Lemma 4.9.** Let  $x, y \in V$ . Then

- (i)  $[x, y] \subseteq \omega$  implies that  $[x, y] \not\subseteq g\omega$ .
- (ii)  $\tau_P \subseteq \Delta_P^g$ .

*Proof.* (i) If  $[x, y] \subseteq \omega$  and  $[x, y] \subseteq g\omega$ , then  $\bar{e} = [\bar{x}, \bar{y}] = \bar{g}$  and so  $g \in \omega$ , a contradiction. Therefore,  $[x, y] \not\subseteq g\omega$ .

- (ii) Clearly, the vertex set of  $\tau_P$  and  $\Delta_P^g$  is  $P - Z(P)$ . Now, if  $\{x, y\}$  is an edge in  $\tau_P$ , then  $xy\omega = yx\omega$ , i.e.  $[x, y] \subseteq \omega$ . It follows from (i), that  $\{x, y\}$  is an edge in  $\Delta_P^g$ . Consequently,  $\tau_P \subseteq \Delta_P^g$ . □



**Lemma 4.10.** *Assume  $K(P) = \{\omega, g\omega\}$  or  $\{\omega, g\omega, g^{-1}\omega\}$ . Then  $\Delta_P^g$  is equal to  $\tau_P$ .*

*Proof.* Obviously, the vertex set of  $\Delta_P^g$  and  $\tau_P$  is  $P - Z(P)$ . Now, let  $\{x, y\}$  is an edge in  $\Delta_P^g$ . Then  $[x, y] \not\subseteq g\omega$ . Using hypotheses and  $[x, y] \in K(P)$ , we get  $[x, y] = \omega$  i.e.  $\{x, y\}$  is an edge in  $\tau_P$ . Similarly, consider  $K(P) = \{\omega, g\omega, g^{-1}\omega\}$  and  $\{x, y\}$  be an edge in  $\Delta_P^g$ . Now, since  $[x, y] \in K(P)$  by  $[x, y] \not\subseteq g\omega$  and  $g^{-1}\omega$ , we conclude that  $[x, y] = \omega$ . Moreover,  $\{x, y\}$  is an edge in  $\tau_P$ . Consequently,  $\Delta_P^g$  is the commuting graph  $\tau_P$ .  $\square$

**Theorem 4.11.** *Consider  $g$  is a non-central element of  $P$ .*

- (i) *If  $g^2 \subseteq \omega$ , then  $\text{diam}(\Delta_P^g) = 2$ .*
- (ii) *If  $g^2 \not\subseteq \omega$  and  $g^3 \not\subseteq \omega$ , then  $\text{diam}(\Delta_P^g) \leq 3$ .*

*Proof.* (i) Take  $x \in P - Z(P)$  such that  $x \neq g$ . If  $[x, g] \subseteq g\omega$ , then  $[\bar{x}, \bar{g}] = \bar{g}$ . It implies that  $\bar{g} = \bar{e}$  and so  $g \in \omega$ , a contradiction. Therefore,

$$(1) \quad [x, g] \not\subseteq g\omega.$$

Also, by  $g^2 \subseteq \omega$ , we get  $\bar{g}^2 = \bar{e}$ , i.e.  $\bar{g}^{-1} = \bar{g}$ . Now, if  $[x, g] \subseteq g^{-1}\omega$ , then  $[\bar{x}, \bar{g}] = \bar{g}^{-1} = \bar{g}$ . It implies that  $\bar{g} = \bar{e}$ , i.e.  $g \in \omega$ , a contradiction. Consequently,

$$(2) \quad [x, g] \not\subseteq g^{-1}\omega.$$

By (1) and (2), we have  $\{x, g\}$  is an edge in  $\Delta_P^g$  and  $\text{diam}(\Delta_P^g) \leq 2$ . On the other hand by Remark 4.6, there exist  $x, y \in P$  such that  $[x, y] \subseteq g\omega, g^{-1}\omega$ . Therefore,  $d(x, y) \geq 2$ . Consequently,  $\text{diam}(\Delta_P^g) = 2$ .

- (ii) We consider two cases, (1)  $[x, g] \not\subseteq g^{-1}\omega$ , (2)  $[x, g] \subseteq g^{-1}\omega$ .  
For (1), we have  $\{x, g\}$  is an edge in  $\Delta_P^g$ .

For (2), by (1), we get

$$(3) \quad \begin{aligned} [x, g^2] &\subseteq [x, g][x, g]^g \subseteq [x, g]g^{-1}[x, g]g \\ &\subseteq [x, g]g^{-1}\omega[x, g]g\omega \\ &\subseteq g^{-1}\omega g^{-1}\omega g^{-1}\omega g\omega. \end{aligned}$$

Since  $g^{-1}g \subseteq \omega$ , we get that  $g^{-1}\omega = g\omega = \omega$ . Then by (3),  $[x, g^2] \subseteq g^{-2}\omega$ . It follows by hypotheses that  $[x, g^2] \subseteq g^{-2}\omega \not\subseteq g\omega$ , and  $g^{-1}\omega$ . Now, for any  $z \in g^2$ ,  $[x, z] \subseteq [x, g^2] \not\subseteq g\omega$  and  $g^{-1}\omega$  and so  $\{x, z\}$  is an edge. Therefore, for any  $x \in P - Z(P)$ , we have  $\{x, g\}$  or  $\{x, z\}$  is an edge, where  $z \in g^2$ . Moreover, for any  $x, y \in V$  if  $x, y$ , join with  $g$  or  $z$ , then  $d(x, y) \leq 2$ . Otherwise  $\{x, g\}$  and  $\{y, z\}$  or  $\{y, g\}$  and  $\{x, z\}$  are edges and so by  $[g, g^2] = g^{-3}g^3 \subseteq \omega$  and Lemma 4.9, we have  $[g, g^2] \not\subseteq g\omega, g^{-1}\omega$ . Therefore,  $d(x, y) \leq 3$ . Hence,  $\text{diam}(\Delta_P^g) \leq 3$ .  $\square$

**Theorem 4.12.** *Let  $|Z(P)| = 1$ ,  $g \in P$ ,  $g \notin \omega$  and  $|C_P(g)| \neq 3$ . Then  $\Delta_P^g$  is connected.*

*Proof.* First, consider  $g^3 \notin \omega$ . Then by Theorem 4.11  $\Delta_P^g$  is connected. Now, if  $g^3 \subseteq \omega$ , then  $g^2 \notin \omega$ . Since  $|C_P(g)| \neq 3$ , we conclude that  $|C_P(g)| > 3$ . Therefore there exist  $a \in C_P(g)$  such that  $a \neq e, g$  and  $g^{-1}$ . We can take  $x_0 \in P - Z(P)$  such that  $[x_0, g^2] \subseteq g\omega$  and  $[x_0, g] \subseteq g^{-1}\omega$ . By Lemma 3.1,  $a \in C_P(g)$  and

$$g\omega = a^{-1}ag\omega = a^{-1}gaw = a^{-1}gaw = g\omega,$$

we have

$$[x_0, ga] \subseteq [x_0, a][x_0, g]^a \subseteq [x_0, a]g^{-a}\omega$$

and

$$\begin{aligned} [x_0, g^2a] &\subseteq [x_0, a][x_0, g^2]^a \subseteq [x_0, a](g^a\omega) \\ &= [x_0, a]g\omega = [x_0, a]g^{-2}\omega. \end{aligned}$$

Since  $a \in C_P(g)$  we have  $[a, g] \subseteq \omega$  and so by Lemma 4.9,  $[a, g] \not\subseteq g^{-1}\omega$ ,  $g\omega$ . Then  $\{a, g\}$  is an edge. Now, if  $[x_0, a] \not\subseteq g\omega$ ,  $g^{-1}\omega$ , then  $\{a, x_0\}$  and  $\{a, g\}$  are edges. Thus,  $\Delta_P^g$  is connected. Otherwise,  $[x_0, a] \subseteq g\omega$  or  $[x_0, a] \subseteq g^{-1}\omega$ . But  $[x_0, a] \subseteq g\omega$ , implies that

$$(4) \quad [x_0, ga] \subseteq [x_0, a]g^{-a}\omega \subseteq g\omega g^{-a}\omega = [g, a]\omega \subseteq \omega$$

We consider two cases (i)  $ga \subseteq Z(P)$ , (ii)  $ga \not\subseteq Z(P)$ .

For (i), by  $|Z(P)| = 1$ , we get  $ga = \{e\}$  and so  $g^{-1} = a$ , a contradiction.

For (ii), we get  $ga \subseteq P - Z(P)$  and so for any  $t \in ga$ , by Theorem 3.1, and hypotheses, we get

$$(5) \quad [x_0, g^2a] \subseteq [x_0, a]g^{-2}\omega \subseteq g^{-3}\omega \subseteq \omega.$$

Again we consider two cases (i)  $g^2a \subseteq Z(P)$ , (ii)  $g^2a \not\subseteq Z(P)$ .

For (i), by hypotheses, we have  $a = g^{-2} = g$ , a contradiction.

For (ii), we have  $g^2a \subseteq P - Z(P)$  and so for any  $t \in g^2a$  by (5), we obtain  $\{t, x_0\}$  and  $\{t, g\}$  are edges, i.e,  $\Delta_P^g$  is connected.  $\square$

*Remark 4.13.* The converse of Theorem 4.12, is not true, because by Example 4.3, we have a connected graph that  $|Z(P)| \neq 1$  and  $C_P(b) = 3$ .

**Theorem 4.14.** *Assume  $P$  is a non-commutative simple polygroup and for any  $x \in P$ ,  $|C_P(x)| \geq 3$ . Then  $\Delta_P^g$  has no isolated vertices.*

*Proof.* Take  $x \in P - Z(P)$ . If  $x^2 \notin \omega$ , then  $[x, x^{-1}] \subseteq \omega$  and so by Lemma 4.9,  $\{x, x^{-1}\}$  is an edge. If  $x^2 \notin \omega$  from  $|C_P(x)| \geq 3$ , there exist  $t \in C_P(x)$  such that  $t \neq e, x$ . Since  $[t, x] \subseteq \omega$  by Lemma 4.9, we have  $\{t, x\}$  is an edge. Therefore  $\Delta_P^g$  has no isolated vertices.  $\square$

**Definition 4.15.** Let  $P$  and  $H$  be two polygroups. Then  $\Delta_P^g$  and  $\Delta_H^h$  are said to be isomorphic, we denote by  $\Delta_P^g \simeq \Delta_H^h$ , if there is a bijection map  $\varphi : P - Z(P) \rightarrow H - Z(H)$  preserving edges, i.e, for any  $x, y \in P - Z(P)$ ,  $\{x, y\}$  is an edge if and only if  $\{\varphi(x), \varphi(y)\}$  is an edge.

**Example 4.16.** Assume  $P$  is as Example 4.3 and the map  $\varphi : P - Z(P) \rightarrow P - Z(P)$  is defined by  $\varphi(b) = c$ ,  $\varphi(c) = b$  and otherwise  $\varphi(x) = x$ . It is easy to see that  $\Delta_P^b \simeq \Delta_P^c$ .

**Lemma 4.17.** Let  $\varphi : P \rightarrow H$  be an isomorphism of polygroups. Then

- (i)  $\varphi(\omega_P) = \omega_H$ .
- (ii)  $\Delta_P^g \simeq \Delta_H^h$ , where  $h = \varphi(g)$ .

*Proof.* (i) Take  $x \in \omega_H$ . Then  $\bar{x} = \bar{e}_H$ , i.e,  $\{x, e_H\} \subseteq \prod_{i=1}^n z_i$  for some  $z_i \in H$ ,  $1 \leq i \leq n$ . Since  $e_H$  and  $x \in \prod_{i=1}^n z_i$ , we have  $\varphi^{-1}(e_H)$  and  $\varphi^{-1}(x) \in \prod_{i=1}^n \varphi^{-1}(z_i)$  and so  $\{\varphi^{-1}(x), \varphi^{-1}(e_H)\} \in \prod_{i=1}^n \varphi^{-1}(z_i)$ . This implies that  $\varphi^{-1}(x) \in \omega_P$ , i.e,  $x \in \varphi(\omega_P)$ . Similarly,  $\varphi(\omega_P) \subseteq \omega_H$ . Therefore,  $\varphi(\omega_P) = \omega_H$ .

- (ii) It is clear by part (i). □

**Lemma 4.18.** Let  $A$  and  $B$  be two polygroups and  $f : A \rightarrow B$  be an isomorphism. Then for any  $a_1, a_2 \in A$ , we have  $[a_1, a_2] \notin \omega_A$  if and only if  $[f(a_1), f(a_2)] \notin \omega_B$ .

*Proof.* It is clear by Lemmas 2.7 and 4.17. □

**Theorem 4.19.** Let  $A$  and  $B$  be two non-commutative isomorphic polygroups. If  $\Delta_P^g \simeq \Delta_H^h$ , then  $\Delta_{P \times A}^{(g,a)} \simeq \Delta_{H \times B}^{(h,b)}$ , where  $h = \varphi(g)$  and  $b = f(a)$  for bijection map  $f : A \rightarrow B$ .

*Proof.* Let  $\varphi : P - Z(P) \rightarrow H - Z(H)$  be a graph isomorphism. Then by Lemma 4.17 (i), for any  $x_1, x_2 \in P - Z(P)$ ,  $[x_1, x_2] \notin g\omega_P$  if and only if  $[\varphi(x_1), \varphi(x_2)] \notin \varphi(g)\omega_H$ . Since  $A, B$  are isomorphic, we have a bijection map  $f : A \rightarrow B$ . Now, for  $x_1, x_2 \in P$  and  $a_1, a_2 \in A$  by Lemma 4.17 (i) and 4.18, we have  $\{(x_1, a_1), (x_2, a_2)\}$  is an edge in  $\Delta_{P \times A}^{(g,a)}$  if and only if  $[(x_1, a_1), (x_2, a_2)] \notin (g, a)\omega_{P \times A}$  if and only if  $([x_1, x_2], [a_1, a_2]) \notin (g, a)\omega_{P \times A}$  if and only if  $[x_1, x_2] \notin g\omega_P$  and  $[a_1, a_2] \notin a\omega_A$  if and only if  $[\varphi(x_1), \varphi(x_2)] \notin \varphi(g)\omega_H$  and  $[f(a_1), f(a_2)] \notin f(a)\omega_B$  if and only if  $[\varphi(x_1), \varphi(x_2)] \notin h\omega_H$  and  $[f(a_1), f(a_2)] \notin b\omega_B$  if and only if  $\{\varphi(x_1), \varphi(x_2)\}$  is an edge in  $\Delta_H^h$  and  $\{f(a_1), f(a_2)\}$  is an edge in  $\Delta_B^b$ . Consequently, by Theorem 3.3  $\Delta_{P \times A}^{(g,a)} \stackrel{(\varphi, f)}{\simeq} \Delta_{H \times B}^{(h,b)}$ . □

**Example 4.20.** Let  $\Delta_P^b$  and  $\Delta_P^c$  be as Examples 4.4 and 4.16. Then  $\Delta_P^b \simeq \Delta_P^c$  and  $\Delta_{P \times P}^{(b,f)} \simeq \Delta_{P \times P}^{(c,f)}$ .

**Theorem 4.21.** Assume  $P_H = H \cup \{b\}$  and  $P_G = G \cup \{a\}$ ,  $g \notin \omega_{P_G}$  and  $f(g) \notin \omega_{P_H}$ . Then  $\Delta_{P_G}^g \stackrel{f}{\simeq} \Delta_{P_H}^{f(g)}$  if and only if  $\Delta_G^g \stackrel{f}{\simeq} \Delta_H^{f(g)}$

*Proof.* ( $\Rightarrow$ ) Let  $g_1, g_2 \in G$ .  $Z(G)$  and  $[g_1, g_2] \neq g$ . Then by Theorem 2.11  $[g_1, g_2] \notin g\omega_{P_G} = \{g\}$ . Since  $\Delta_{P_G}^g \stackrel{f}{\simeq} \Delta_{P_H}^{f(g)}$ , there exists a bijection  $f : P_G - Z(P_G) \rightarrow P_H - Z(P_H)$  and  $[f(g_1), f(g_2)] \notin f(g)\omega_{P_H} = \{f(g)\}$ . It follows that  $[f(g_1), f(g_2)] \neq f(g)$ . Therefore,  $\Delta_G^g \stackrel{f}{\simeq} \Delta_H^{f(g)}$ , where  $f|_{G-Z(G)} : G - Z(G) \rightarrow H - Z(H)$ , is a bijection.

( $\Leftarrow$ ) Since  $\Delta_G^g \stackrel{f}{\simeq} \Delta_H^{f(g)}$ , there is a bijection  $f : G - Z(G) \rightarrow H - Z(H)$ . Then  $f : P_G - Z(P_G) \rightarrow P_H - Z(P_H)$  is a bijection. If  $x, y \in P_G - Z(P_G)$  and  $[x, y] \notin g\omega_{P_G} = \{g\}$ , then  $[x, y] \neq g$ . Since  $\Delta_G^g \stackrel{f}{\simeq} \Delta_H^{f(g)}$ , we conclude that  $[f(x), f(y)] \neq f(g)$  and so  $[f(x), f(y)] \notin f(g)\omega_{P_H} = \{f(g)\}$ . Consequently,  $\Delta_{P_G}^g \stackrel{f}{\simeq} \Delta_{P_H}^{f(g)}$ .  $\square$

**Definition 4.22.** [6] A polygroup  $(P, \circ)$  is called a polygroup of *exponent*  $n$  ( $n \in \mathbb{N}$ ) if for each non-trivial element  $x$  of  $P$  we have  $\underbrace{x \circ x \circ \dots \circ x}_n = e$ .

**Theorem 4.23.** Every polygroup of exponent 3 is a group. In addition, if  $P$  and  $H$  are two finite non-abelian polygroups of exponent 3 and  $\Delta_P^g \simeq \Delta_H^h$ , for some non-identity element  $h \in H$ . Then  $|Z(P)| = |Z(H)|$ .

*Proof.* Let  $P$  be a polygroup of exponent 3 and  $b \in P$  be an arbitrary element. Since  $bbb = e$ , then  $bx = e$  for some  $x \in bb$ . Thus  $b^{-1} = x$ , which implies

$$bb^{-1} = e \text{ for any } b \in P. (*)$$

Let  $x \in yz$ . By Definition 2.1, we have  $z \in y^{-1}x$ . Thus by (\*)  $x \in yz \subseteq yy^{-1}x = x$ . Therefore,  $x = yz$  and so every polygroups of exponent 3 is a group.

In addition, consider  $P$  and  $H$  are two polygroups of exponent 3 and  $\Delta_P^g \simeq \Delta_H^h$ . Then  $|Z(P)| = |Z(H)|$ .  $\square$

**Theorem 4.24.** [16] Assume  $G$  is a finite nilpotent non-abelian group of add order and  $\Delta_G^g$  has no vertex adjacent to all other vertices. If  $\Delta_G^g \simeq \Delta_H^h$ , then  $H$  is nilpotent

**Theorem 4.25.** Assume  $G$  is a finite non-abelian group of add order and  $\Delta_G^g$  has no vertex adjacent to all other vertices. If  $\Delta_{P_G}^g \stackrel{f}{\simeq} \Delta_{P_H}^{f(g)}$  and  $P_G$  is nilpotent, then  $P_H$  is nilpotent.

*Proof.* Since  $\Delta_{P_G}^g \stackrel{f}{\simeq} \Delta_{P_H}^{f(g)}$  by 4.21, we conclude that  $\Delta_G^g \stackrel{f}{\simeq} \Delta_H^{f(g)}$ . Also, by Theorem 3.4,  $G$  is nilpotent. It follows by Theorem 4.24, that  $H$  is nilpotent. Consequently, by Theorem 3.4,  $P_H$  is nilpotent.  $\square$

**Corollary 4.26.** *Let  $G$  and  $H$  be two non-abelian finite groups such that  $\Delta_{P_G}^g \stackrel{f}{\simeq} \Delta_{P_H}^{f(g)}$ , for some non-identity element  $h \in H$ . Then  $|P_G| = |P_H|$ .*

*Proof.* Since  $\Delta_{P_G}^g \stackrel{f}{\simeq} \Delta_{P_H}^{f(g)}$ , by Theorem 4.21, we get  $\Delta_G^g \simeq \Delta_H^{f(g)}$ . Thus by Theorem 4.23,  $|G| = |H|$ . Consequently  $|P_G| = |P_H|$ .  $\square$

## 5. Conclusion

*In this paper, some results on the commutator of polygroups were obtained. Moreover, a relation between nilpotent polygroups and groups was obtained. Then the notion of  $g$ -graph was introduced. Also, some properties of this graph, such as diameter and connectivity are stated. Moreover, isomorphic  $g$ -graphs were investigated. This paper would be useful for studying nilpotent polygroups. In addition, you can obtain some results on Engel and solvable polygroups by the relation between these structures. Also, by this method, you can study other algebraic structures such as hypergroups, hyperrings, and so on.*

## References

- [1] Abdollahi, A., & Akbari, S., & Maimani, H. R. (2006). Non-commuting graph of a group. *Journal of Algebra*, 298, 468-492. <https://doi.org/10.1016/J.JALGEBRA.2006.02.015>.
- [2] Ameri, R., & Mohammadzadeh, E. (2015). Engel groups derived from hypergroups. *European Journal of Combinatorics*, 44, 191-197. <https://doi.org/10.1016/j.ejc.2014.08.004>
- [3] Corsini, P. (1993). *Prolegomena of hypergroup theory*. Aviani Editore, Tricesimo.
- [4] Corsini, P., & V. Leoreanu, V. (2003). *Applications of hyperstructure theory*. Kluwer Academic Publishers, Dordrecht. <https://link.springer.com/book/10.1007/978-1-4757-3714-1>.
- [5] Comer, S. D. (1982). Extension of polygroups by polygroups and their representations using colour schemes. *Lecture notes in Math*, No 1004, Universal Algebra and Lattice Theory, 91-103. <https://link.springer.com/chapter/10.1007/BFb0063431>.
- [6] Davvaz, B. (2013). *Polygroup theory and related systems*. World Scientific. <https://doi.org/10.1142/8593>.
- [7] Freni, D. (1991). Une note sur le cœur dun hypergroup et sur la cloture transitive  $\beta^*$  de  $\beta$  (A note on the core of a hypergroup and the transitive closure  $\beta^*$  of  $\beta$ ). *Riv. Mat. Pura Appl*, 8, 153-156 (in French).
- [8] Hamidi, M., & Borumand Saeid, A. (2017). Accessible single-valued neutrosophic graphs. *Journal of Applied Mathematics and Computing*, 57, 121-146. <https://doi.org/10.1007/s12190-017-1098-z>.
- [9] Hamidi, M., & Mirvakili, S., & Hatami, A. (2022). Interval-valued grey (hyper) group. *Journal of Algebraic Hyperstructures and Logical Algebras*, 3(3), 85-96. <https://10.52547/HATEF.JAHLA.3.3.6>
- [10] Koskas, M., & Groupoides. (1970). demi-hypergroups ehypergroupes. *Journal de mathématiques pures et Appliquées*, 49, 155-192. <https://10.1007/s12190-017-1098-z>.

- [11] Marty, F. (1934). Sur une Generalization de la Notion de Groupe. in: 8th Congress Math. Scandenaves, Stockholm, Sweden, 45-49.
- [12] Moghaddamfar, A. R. (2006). About noncommuting graphs. Siberian Mathematical Journal, 47(5), 911–914. <https://10.1007/s11202-006-0101-y>.
- [13] Moghaddamfar, A. R., & Shi, W. J., & Zhou, W., & Zokayi, A. R. (2005). On the noncommuting graph associated with a finite group. Siberian Math. J., 46(2), 325–332. <https://10.1007/s11202-005-0034-x>.
- [14] Mohammadzadeh, E. (2021). A graph related to a polygroup with respect to automorphism. Journal of Algebraic Hyperstructures and Logical Algebras, 2(1), 99-112. <https://10.52547/HATEF.JAHLA.2.1.7>.
- [15] Mohammadzadeh, F., & Mohammadzadeh, E. (2022). On nilpotent polygroups. 10th national mathematics conference of the Payame Noor University, Shiraz, 1-3.
- [16] Nasiri, M., & Erfanian, A., & Gangali, M., & Jafarzadeh, A. (2017). Isomorphic g-noncommuting graphs of finite groups. Publ. Math. Debrecen, 91/1-2, 33–42. <https://10.5486/pmd.2017.7577>.
- [17] M. Suzuki. (1982). Group theory I. Springer-verlag, New York. <https://www.springer.com/series/1721>.

ELAHE MOHAMMADZADEH  
ORCID NUMBER: 0000-0003-3348-5780  
DEPARTMENT OF MATHEMATICS  
UNIVERSITY OF PAYAME NOOR  
TEHRAN, IRAN

*Email address: mohamadzadeh@pnu.ac.ir*

SOMAYEH AMIRMOFIDI  
ORCID NUMBER: 0009-0002-4308-2758  
DEPARTMENT OF MATHEMATICS  
UNIVERSITY OF PAYAME NOOR  
TEHRAN, IRAN

*Email address: s.amirmofidi@student.pnu.ac.ir*

MOHAMMAD HAMIDI  
ORCID NUMBER: 0000-0002-8686-6942  
DEPARTMENT OF MATHEMATICS  
UNIVERSITY OF PAYAME NOOR  
TEHRAN, IRAN

*Email address: m.hamidi@pnu.ac.ir*